

## **Routine Preventive Pediatric Health Care – Adolescence**

The following guideline provides recommendations for routine preventive services for children 11 years to 20 years. Children at increased risk may warrant additional services.

Age	11	12	13	14	15	16	17	18	19	20
	yrs									
HISTORY Initial/Interval	Х	X	X	X	X	X	Х	X	X	Х
MEASUREMENTS										
Length/Height and Weight	X	Х	Х	X	X	X	Х	X	Х	Χ
Body Mass Index	Х	X	Х	Χ	Х	Х	Х	Х	Х	Χ
Blood Pressure	Х	Х	Х	Х	Х	Х	Х	Х	Х	Χ
SENSORY SCREENING										
Vision	*	Х	*	*	Х	*	*	*	*	*
Hearing Screen with audiometry including 6000 & 8000 Hz high	<	<	Х	>	<	Х	>	<	<	Χ
frequencies once between 11 and 14 years, once between 15 and 17 years, and										
once between 18 and 21 years.										
DEVELOPMENTAL/BEHAVIORAL ASSESSMENT										
Developmental Surveillance	Х	Х	Χ	Χ	Χ	Χ	Χ	Х	Х	Х
Psychosocial/Behavioral Assessment Should be family centered; may	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Х
include assessment of child social-emotional health, caregiver depression, and										
social determinants of health.										
http://pediatrics.aappublications.org/content/135/2/384										
http://pediatrics.aappublications.org.content/137/4/e20160039										
Tobacco, Alcohol, or Drug Use Assessment. A recommended screening tool	*	*	*	*	*	*	*	*	*	*
is available at http://www.ceasar-boston.org/CRAFFT/index.php.										
Depression Screening. Recommended screening using the Patient Health		Х	Х	Χ	Х	Х	Х	Х	Х	Х
Questionnaire (PHQ)-2 or other tools available in the GLAD-PC toolkit and at										
http://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Mental-										
Health/Documents/MH_ScreeningChart.pdf.										
PHYSICAL EXAMINATION	Х	Х	Х	Χ	Х	Х	Х	Х	Х	Χ



Age	11	12	13	14	15	16	17	18	19	20
	yrs									
Procedures										
Immunization Every visit should be an opportunity to update and	Х	Х	Х	Χ	Х	Х	Χ	Х	Х	Х
complete a child's immunizations. Schedules are available at										
http://aapredbook.aappublications.org/site/resources/izschedules.xhtml.										
Anemia	*	*	*	*	*	*	*	*	*	*
Tuberculosis Testing. Tuberculosis testing per recommendations of the	*	*	*	*	*	*	*	*	*	*
Committee on Infectious Diseases, published in the current edition of AAP Red										
Book: Report of the Committee on Infectious Diseases. Testing should be										
performed on recognition of high-risk factors.										
Dyslipidemia Screening	>	*	*	*	*	*	<	<	<	Χ
Sexually Transmitted Infections	*	*	*	*	*	*	*	*	*	*
HIV Once between the ages of 15 & 18, making every effort to preserve	*	*	*	*	<	<	Х	>	*	*
confidentiality of the adolescent.										
Fluoride Supplementation	*	*	*	*	*	*				
ANTICIPATORY GUIDANCE	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х

This guideline is based on the Bright Futures and the American Academy of Pediatrics Recommendations for Preventive Pediatric Health Care. Individual patient considerations and advances in medical science may supersede or modify these recommendations.

July 10, 2018

KEY X = to be performed \* = risk assessment to be performed with appropriate action to follow, if positive <<<X>>> = range during which a service may be provided